High Sensitivity C-Reactive Protein

		Specimen 1			Specimen 2			Specimen 3			Specimen 4			Specimen 5			
Name	**Line No.**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**No. of Labs**
High Sensitivity C-Reactive Protein																	
Initial Grouping by Reagent																	
Abbott Aeroset/Architect	1	0.72 - 1.09	S 0.902	0.061	4.73 - 5.99	S 5.359	0.209	-			-			-			10
Beckman Olympus	2	0.82 - 1.03	S 0.924	0.035	4.17 - 6.13	S 5.149	0.326	-			-			-			13
Siemens Dim CardioPhase	3	0.68 - 1.54	S 1.112	0.144	4.62 - 7.35	S 5.986	0.455	-			-			-			15
Initial Grouping by Sensitivity or Principle																	
All direct quantitive meth	4	0 - 1.81	S 0.863	0.316	3.64 - 7.74	S 5.691	0.682	-			-			-			62
All wide range methods	5	0.15 - 1.56	S 0.857	0.235	4.17 - 6.13	S 5.149	0.326	-			-			-			14
All Roche HS methods	6	0.64 - 1.3	S 0.968	0.111	4.39 - 6.06	S 5.224	0.278	-			-			-			18
Total Population																	
Whole Population	7	0.07 - 1.71	S 0.887	0.273	3.56 - 7.58	S 5.567	0.67	-			-			-			99