Sex Hormone Binding Globulin/Testosterone

		Specimen 1			Specimen 2			Specimen 3			Specimen 4			Specimen 5			
Name	**Line No.**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**No. of Labs**
Sex Hormone Binding Globulin																	
Initial Grouping by Reagent																	
Beckman Coulter Access	1	17.3 - 32.2	P 24.77	1.67	42.5 - 78.9	P 60.72	5.58	-			-			-			10
Initial Grouping by Sensitivity or Principle																	
Luminometric	2	17.9 - 33.3	P 25.64	2.04	43.7 - 81.2	P 62.43	5.93	-			-			-			19
Total Population																	
Whole Population	3	18.1 - 33.6	P 25.87	2.31	43.5 - 80.8	P 62.14	6.65	-			-			-			26
Testosterone																	
Initial Grouping by Sensitivity or Principle																	
Luminometric	1	79 - 146	P 112.2	7.2	933 - 1733	P 1333.2	149.4	-			-			-			15
Total Population																	
Whole Population	2	77 - 144	P 110.5	7.8	962 - 1787	P 1374.3	193.7	-			-			-			21
Testosterone, Bioavailable																	
Total Population																	
Whole Population	1	43.6 - 81.0	P 62.34	14.92	333.7 - 619.7	P 476.67	40.66	-			-			-			8
Testosterone, Free																	
Initial Grouping by Reagent																	
Albumin equal to 4.3 mg/dL	1	1.7 - 3.1	P 2.39	0.16	15.9 - 29.5	P 22.72	3.21	-			-			-			11
Initial Grouping by Sensitivity or Principle																	
All other methods	2	1.7 - 3.1	P 2.39	0.16	15.9 - 29.5	P 22.72	3.21	-			-			-			11
Total Population																	
Whole Population	3	1.7 - 3.1	P 2.39	0.16	15.9 - 29.5	P 22.72	3.21	-			-			-			11