High Sensitivity C-Reactive Protein

		Specimen 1			Specimen 2			Specimen 3			Specimen 4			Specimen 5			
Name	**Line No.**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**No. of Labs**
High Sensitivity C-Reactive Protein																	
Initial Grouping by Reagent																	
Beckman Olympus	1	2.71 - 3.98 S	3.343	0.211	0.95 - 1.31 S	1.131	0.061	-			-			-			17
Siemens Dim CardioPhase	2	3.78 - 4.45 S	4.116	0.113	1.22 - 1.54 S	1.383	0.053	-			-			-			13
Initial Grouping by Sensitivity or Principle																	
All direct quantitive meth	3	0.33 - 6.88 S	3.604	1.093	0 - 2.19 S	1.035	0.384	-			-			-			54
All wide range methods	4	1.31 - 5.08 S	3.195	0.628	0.58 - 1.61 S	1.092	0.171	-			-			-			18
All Roche HS methods	5	2.59 - 4.28 S	3.436	0.281	0.7 - 1.66 S	1.179	0.16	-			-			-			15
Total Population																	
Whole Population	6	0.82 - 6.23 S	3.524	0.901	0.14 - 2.03 S	1.085	0.314	-			-			-			94