Sex Hormone Binding Globulin/Testosterone

		Specimen 1			Specimen 2			Specimen 3			Specimen 4			Specimen 5			
Name	**Line No.**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**No. of Labs**
Sex Hormone Binding Globulin																	
Initial Grouping by Reagent																	
Beckman Coulter Access	1	27.1 - 50.3	P 38.69	3.97	23.6 - 43.8	P 33.69	2.46	-			-			-			11
Initial Grouping by Sensitivity or Principle																	
Luminometric	2	28.0 - 52.0	P 40.0	4.42	23.6 - 43.9	P 33.75	2.82	-			-			-			20
Total Population																	
Whole Population	3	27.6 - 51.2	P 39.39	4.51	23.4 - 43.4	P 33.41	3.14	-			-			-			26
Testosterone																	
Initial Grouping by Sensitivity or Principle																	
Luminometric	1	515 - 957	P 736.0	138.5	337 - 626	P 481.6	99.3	-			-			-			15
Total Population																	
Whole Population	2	522 - 970	P 746.0	156.7	342 - 635	P 488.5	106.2	-			-			-			20
Testosterone, Bioavailable																	
Total Population																	
Whole Population	1	255.2 - 473.9	P 364.5	64.46	157.3 - 292.2	P 224.77	20.81	-			-			-			8
Testosterone, Free																	
Initial Grouping by Reagent																	
Albumin equal to 4.3 mg/dL	1	11.5 - 21.4	P 16.48	2.59	6.9 - 12.7	P 9.8	1.12	-			-			-			10
Initial Grouping by Sensitivity or Principle																	
All other methods	2	11.5 - 21.4	P 16.48	2.59	6.9 - 12.7	P 9.8	1.12	-			-			-			10
Total Population																	
Whole Population	3	11.5 - 21.4	P 16.44	2.46	6.9 - 12.8	P 9.82	1.06	-			-			-			11