High Sensitivity C-Reactive Protein

		Specimen 1			Specimen 2			Specimen 3			Specimen 4			Specimen 5			
Name	**Line No.**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**No. of Labs**
High Sensitivity C-Reactive Protein																	
Initial Grouping by Reagent																	
Abbott Aeroset/Architect	1	1.93 - 17.02 S	9.475	2.516	1.92 - 2.44 S	2.178	0.087	-			-			-			10
Beckman Olympus	2	9.16 - 11.08 S	10.122	0.32	1.91 - 2.39 S	2.151	0.081	-			-			-			14
Siemens Dim CardioPhase	3	8.41 - 11.35 S	9.877	0.489	1.93 - 3.46 S	2.696	0.256	-			-			-			15
Initial Grouping by Sensitivity or Principle																	
All direct quantitive meth	4	4.54 - 15.67 S	10.106	1.855	0.32 - 4.05 S	2.188	0.622	-			-			-			63
All wide range methods	5	9.16 - 11.08 S	10.122	0.32	0.55 - 3.49 S	2.022	0.491	-			-			-			15
All Roche HS methods	6	8.77 - 11.77 S	10.266	0.5	1.75 - 3.08 S	2.419	0.222	-			-			-			17
Total Population																	
Whole Population	7	5.64 - 14.73 S	10.181	1.515	0.57 - 3.87 S	2.219	0.55	-			-			-			101