

Prostate Specific Ag, Free**Initial Grouping by Reagent and Instrument**

Beckman Coulter Access & Beckman Coulter Access	1	2.7 - 5.0	P 3.83	0.33	4.1 - 7.7	P 5.89	0.5	-	-	-	11
---	---	-----------	--------	------	-----------	--------	-----	---	---	---	----

Initial Grouping by Reagent

Beckman Coulter Access	2	2.7 - 5.1	P 3.89	0.28	4.1 - 7.6	P 5.87	0.43	-	-	-	17
Roche Elecsys	3	1.8 - 3.6	C 2.72	0.09	2.9 - 5.4	P 4.13	0.14	-	-	-	11

Initial Grouping by Sensitivity or Principle

Moderate recovery methods	4	2.4 - 4.4	P 3.41	0.6	3.6 - 6.7	P 5.17	0.9	-	-	-	30
High recovery methods	5	1.8 - 3.6	C 2.72	0.09	2.9 - 5.4	P 4.13	0.14	-	-	-	11

Total Population

Whole Population	6	2.3 - 4.2	P 3.22	0.6	3.4 - 6.3	P 4.88	0.9	-	-	-	41
------------------	---	-----------	--------	-----	-----------	--------	-----	---	---	---	----

Prostatic Acid Phosphatase (PAP)**Total Population**

Whole Population	1	10.5 - 19.4	P 14.95	0.35	10.5 - 19.5	P 15.0	0.5	-	-	-	2
------------------	---	-------------	---------	------	-------------	--------	-----	---	---	---	---

Thyroglobulin**Total Population**

Whole Population	1	6.3 - 27.4	S 16.86	3.52	0 - 75.7	S 29.47	15.4	-	-	-	7
------------------	---	------------	---------	------	----------	---------	------	---	---	---	---