High Sensitivity C-Reactive Protein

		Specimen 1			Specimen 2			Specimen 3			Specimen 4			Specimen 5			
Name	**Line No.**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**Range & Type**	**Mean**	**SD**	**No. of Labs**
High Sensitivity C-Reactive Protein																	
Initial Grouping by Reagent																	
Beckman Olympus	1	2.61 - 3.5	S 3.056	0.149	0.96 - 1.24	S 1.101	0.046	-			-			-			18
Siemens Dim CardioPhase	2	3.39 - 4.26	S 3.825	0.146	1.11 - 1.62	S 1.362	0.085	-			-			-			14
Abbott Aeroset/Architect	3	1.05 - 4.76	S 2.906	0.618	0 - 3.06	S 1.274	0.597	-			-			-			10
Initial Grouping by Sensitivity or Principle																	
All wide range methods	4	1.06 - 4.61	S 2.835	0.591	0.09 - 1.91	S 0.999	0.302	-			-			-			21
All direct quantitive meth	5	1.11 - 5.66	S 3.384	0.759	0 - 3.26	S 1.219	0.68	-			-			-			61
All Roche HS methods	6	2.55 - 3.78	S 3.166	0.204	0.65 - 1.6	S 1.127	0.159	-			-			-			15
Total Population																	
Whole Population	7	1.19 - 5.33	S 3.257	0.691	0 - 2.81	S 1.159	0.551	-			-			-			103